



Globally Rare Species



Earleaf Foxglove (*Agalinis auriculata*)

Habitat: Wet-mesic prairie
Best ID Time: Late-August through early-September






Forked Aster (*Eurybia furcata*)

Habitat: Dry-mesic to mesic hardwoods, often adjacent to lakes or streams, or on slopes with dolomite near the surface
Best ID Time: Late-August through late-September

Clustered Poppy Mallow (*Callirhoe triangulata*)

Habitat: Sand terrace prairies
Best ID Time: Early-July through September






Hill's Thistle (*Cirsium hillii*)

Habitat: Dry prairies and oak barrens
Best ID Time: Late-May through early-June

Ram's Head Lady's-slipper (*Cypripedium arietinum*)

Habitat: Two habitats: cool, dense, calcareous conifer swamps and cool, sandy margins of conifer forests, boreal forests, and dunes
Best ID Time: Late-May through early-June



Kitten Tails (*Synthyris bullii*)

Habitat: Small woodland openings or near bluff edges
Best ID Time: Late-May through August

Thickspike (*Elymus lanceolatus*)

Habitat: Dunes
Best ID Time: Early-June through August






Western Jacob's Ladder (*Polemonium occidentale ssp. lacustre*)

Habitat: Openings in tamarack swamps
Best ID Time: July

Cliff Cudweed (*Pseudognaphalium saxicola*)

Habitat: Dry to moist, shaded sandstone ledges along the Kickapoo and Wisconsin Rivers
Best ID Time: Early-August through early-October



Goblin Moonwort (*Botrychium mormo*)

Habitat: Northern mesic forest with unwormed soil
Best ID Time: July-September